

## DOCUMENT RESUME

ED 223 666

TM 820 790

AUTHOR Lord, Frederic M.; Wingersky, Marilyn S.  
TITLE Sampling Variances and Covariances of Parameter Estimates in Item Response Theory.  
INSTITUTION Educational Testing Service, Princeton, N.J.  
SPONS AGENCY Office of Naval Research, Arlington, Va. Personnel and Training Research Programs Office.  
REPORT NO ETS-RR-82-33-ONR  
PUB DATE Aug 82  
CONTRACT N00014-80-C-0402  
NOTE 5lp.  
PUB TYPE Reports - Research/Technical (143)

EDRS PRICE MF01/PC03 Plus Postage.  
DESCRIPTORS \*Error of Measurement; \*Estimation (Mathematics); \*Latent Trait Theory; Matrices; \*Maximum Likelihood Statistics; Statistics  
IDENTIFIERS \*Item Parameters

## ABSTRACT

A possible method is developed for computing the asymptotic sampling variance-covariance matrix of joint maximum likelihood estimates in item response theory when both item parameters and abilities are unknown. For a set of artificial data, results are compared with empirical values and with the variance-covariance matrices found by the usual formulas for the case where the abilities are known, or where the item parameters are known. The results are consistent with the conjecture that the new method is asymptotically correct except for errors due to grouping.  
(Author/PN)

\*\*\*\*\*  
\* Reproductions supplied by EDRS are the best that can be made \*  
\* from the original document. \*  
\*\*\*\*\*

U.S. DEPARTMENT OF EDUCATION  
NATIONAL INSTITUTE OF EDUCATION  
EDUCATIONAL RESOURCES INFORMATION  
CENTER (ERIC)

☒ This document has been reproduced as received from the person or organization originating it.

☐ Minor changes have been made to improve reproduction quality.

• Points of view or opinions stated in this document do not necessarily represent official NIE position or policy.

# SAMPLING VARIANCES AND COVARIANCES OF PARAMETER ESTIMATES IN ITEM RESPONSE THEORY

Frederic M. Lord  
and  
Marilyn S. Wingersky

"PERMISSION TO REPRODUCE THIS  
MATERIAL HAS BEEN GRANTED BY

the Office of  
Naval Research

TO THE EDUCATIONAL RESOURCES  
INFORMATION CENTER (ERIC)."

This research was sponsored in part by the  
Personnel and Training Research Programs  
Psychological Sciences Division  
Office of Naval Research, under  
Contract No. N00014-80-C-0402

Contract Authority Identification Number  
NR No. 150-453

Frederic M. Lord, Principal Investigator



Educational Testing Service  
Princeton, New Jersey

August 1982

Reproduction in whole or in part is permitted  
for any purpose of the United States Government.

Approved for public release; distribution  
unlimited.

ED223666

TM 820 792

SAMPLING VARIANCES AND COVARIANCES  
OF PARAMETER ESTIMATES IN  
ITEM RESPONSE THEORY

Frederic M. Lord  
and  
Marilyn S. Wingersky

This research was sponsored in part by the  
Personnel and Training Research Programs  
Psychological Sciences Division  
Office of Naval Research, under  
Contract No. N00014-80-C-0402

Contract Authority Identification Number  
NR No. 150-453

Frederic M. Lord, Principal Investigator

Educational Testing Service  
Princeton, New Jersey

August 1982

Reproduction in whole or in part is permitted  
for any purpose of the United States Government.

Approved for public release; distribution  
unlimited.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS / BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Sampling Variances and Covariances of Parameter Estimates in Item Response Theory		5. TYPE OF REPORT & PERIOD COVERED Technical Report
7. AUTHOR(s) Frédéric M. Lord and Marilyn S. Wingersky		6. PERFORMING ORG. REPORT NUMBER RR-82-33-ONR
9. PERFORMING ORGANIZATION NAME AND ADDRESS Educational Testing Service Princeton, NJ 08541		8. CONTRACT OR GRANT NUMBER(s) N00014-80-C-0402
11. CONTROLLING OFFICE NAME AND ADDRESS Personnel and Training Research Programs Office of Naval Research (Code 458) Arlington, VA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 150-453
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE August 1982
		13. NUMBER OF PAGES 36
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Sampling Covariances                      Item Parameters Standard Errors                              Matrix Inversion Item Response Theory Maximum Likelihood Estimators		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This paper develops a possible method for computing the asymptotic sampling variance-covariance matrix of joint maximum likelihood estimates in item response theory when both item parameters and abilities are unknown. For a set of artificial data, results are compared with empirical values; also with the variance-covariance matrices found by the usual formulas for the case where the abilities are known, or where the item parameters are known. The results are consistent with the conjecture that the new method is asymptotically correct except for errors due to grouping.		

DD FORM 1473

EDITION OF 1 NOV 65 IS OBSOLETE

N 0102-1E (14-6001)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Sampling Variances and Covariances  
of Parameter Estimates in Item Response Theory

Abstract

This paper develops a possible method for computing the asymptotic sampling variance-covariance matrix of joint maximum likelihood estimates in item response theory when both item parameters and abilities are unknown. For a set of artificial data, results are compared with empirical values; also with the variance-covariance matrices found by the usual formulas for the case where the abilities are known, or where the item parameters are known. The results are consistent with the conjecture that the new method is asymptotically correct except for errors due to grouping.

Sampling Variances and Covariances  
of Parameter Estimates in Item Response Theory\*

In item response theory (IRT), the observations come in the form of an  $n$ -by- $N$  matrix, with one row for each item and one column for each examinee. The joint frequency distribution of the observations depends on a vector of  $N$  'ability' parameters--one for each person--and on a matrix of item parameters. Here, we will consider only the three-parameter logistic model for dichotomously scored items, so there will be three item parameters ( $a$ ,  $b$ , and  $c$ ) for each of  $n$  items. A method will be developed for computing the asymptotic sampling variance-covariance matrix when both abilities and item parameters are unknown. Until this is done we do not know the standard errors of the parameter estimates, which handicaps development of a goodness-of fit test and other statistics required in applications of IRT.

If the item (ability) parameters are known, the estimated ability (item) parameters have independent sampling distributions. It can be shown (see Bradley & Gart, 1962) that the maximum likelihood estimates of the ability (item) parameters are consistent. Hence the asymptotic sampling variance for an estimated ability parameter is given by the usual formula

$$\text{Var}(\hat{\tau}_r | a, b, c) = [\mathcal{E}(\partial \ell / \partial \tau_r)^2]^{-1}, \quad (1a)$$

where  $\hat{\tau}_r$  is the estimated ability parameter,  $\ell$  is the log of the likelihood, and  $a$ ,  $b$ , and  $c$  are the known vectors of item parameters.

---

\*This work was supported in part by contract N00014-80-C-0402, project designation NR 150-453 between the Office of Naval Research and Educational Testing Service. Reproduction in whole or in part is permitted for any purpose of the United States Government.

Similarly the asymptotic sampling variance-covariance matrix of the estimated item parameters for an item is given by

$$\| \text{Cov}(\hat{\tau}_v, \hat{\tau}_w | \theta) \| = \| \mathcal{E} \left( \frac{\partial \ell}{\partial \tau_v} \frac{\partial \ell}{\partial \tau_w} \right) \|^{-1} \quad (v, w = 1, 2, 3) \quad (1b)$$

where  $\{\hat{\tau}_v\}$  is a vector consisting of the estimated  $a$ ,  $b$ , and  $c$  for a single item and  $\theta$  is the known vector of abilities.

The right-hand side is the inverse of a 3-by-3 matrix.

When neither item nor ability parameters are known, all parameters are often estimated simultaneously by maximum likelihood. In the (Rasch) case where there is only one parameter per item, Haberman (1977) has shown that all parameter estimates will converge to their true values (will be consistent) when the number of examinees and the number of test items become large simultaneously. Empirical results suggest that consistency probably also holds when all parameters are estimated simultaneously under the three-parameter model. If so, it is reasonable that the asymptotic sampling variance-covariance matrix of all estimated parameters will be given by the usual formula

$$\| \text{Cov}(\hat{\tau}_p, \hat{\tau}_q) \| = \| \mathcal{E} \left( \frac{\partial \ell}{\partial \tau_p} \frac{\partial \ell}{\partial \tau_q} \right) \|^{-1} \quad (p, q = 1, 2, \dots, M) \quad (2)$$

where  $M = 3n + N - 2$  and  $\tau \equiv \{\tau_p\} \equiv \{a_1, b_1, c_1, a_2, b_2, c_2, \dots, a_n, b_n, c_n; \theta_1, \theta_2, \dots, \theta_{N-2}\}$ .

Since standard errors are urgently needed in practical work where all parameters are estimated simultaneously by maximum likelihood, this report compares numerical values provided by (2) with values provided by (1) and with empirically observed sampling fluctuations. The comparisons to be presented suggest that (2) provides useful values for the desired standard errors.

There are several special problems that arise in the evaluation and practical utilization of (2), problems that do not arise in the situation where (1) is appropriate:

1. Until an origin and scale are specified, the parameters are not identifiable.
2. The mathematical formulation is complicated by the choice of origin and scale.
3. The usual choice of origin and scale when estimating IRT parameters is inconvenient for mathematical purposes.
4. The numerical values of the sampling variances are very much affected by the choice of origin and scale.
5. Equation (2) requires the inversion of a matrix of order  $N + 3n - 2$  where  $N$  may be several thousand.

These problems will be considered in subsequent sections.

### 1. Parameterization

The appropriate likelihood function is (Lord, 1980)

$$L(a, b, c; \theta | U) = \prod_{i=1}^n \prod_{a=1}^N \frac{u_{ia}^{1-u_{ia}}}{p_{ia}^{u_{ia}} q_{ia}^{1-u_{ia}}} \quad (3)$$



where  $\theta$  is the vector of the  $N$  ability parameters;  $a$ ,  $b$ , and  $c$  are each a vector of  $n$  item parameters,  $U \equiv \|u_{ia}\|$  is the matrix of item responses  $u_{ia}$  ( $= 0$  or  $1$ ); finally  $Q_{ia} \equiv 1 - P_{ia}$  and  $P_{ia}$  is the item response function; the probability of a correct answer by examinee  $a$  to item  $i$ . Each given  $P_{ia}$  is a function of  $\theta_a$  and of  $a_i$ ,  $b_i$ , and  $c_i$ , but not of any other parameters. In numerical work here,  $P_{ia}$  will be taken to be the three-parameter logistic function

$$P_{ia} \equiv c_i + \frac{1 - c_i}{1 + \exp[-1.7a_i(\theta_a - b_i)]} \quad (4)$$

For mathematical purposes, however, it is only necessary to state that  $P_{ia}$  is an increasing function of  $\theta_a$ .

If we add some constant to all  $\theta_a$  and subtract the same constant from all  $b_i$ , all  $P_{ia}$  will be unchanged. This means that the origin used for measuring ability is entirely arbitrary. If we multiply each  $\theta_a$  and each  $b_i$  by some constant and divide each  $a_i$  by the same constant, again all  $P_{ia}$  will be unchanged. This means that the unit used to measure ability is entirely arbitrary. Since we can change the origin and unit of the  $\theta_a$  without changing (3), it follows that  $\theta$ ,  $a$ ,  $b$ , and  $c$  are not identifiable and cannot be estimated from (3) without further specification.

To conform to a commonly used procedure, we could choose the origin and scale so that for some specified group of examinees the mean of the  $\theta_a$  is zero and the variance is one. This is not convenient mathematically, however. Instead, two other methods of

specifying the origin and scale will be used, even though this will complicate matters later on when the results are applied in practice. In the first method, without loss of generality, arbitrary numerical values will be assigned to  $\theta_{N-1}$  and to  $\theta_N$ .

The  $M \equiv N + 3n - 2$  likelihood equations are

$$0 = \sum_{i=1}^n \sum_{a=1}^N (u_{ia} - P_{ia}) \frac{P_{ia}}{P_{ia} Q_{ia}} \quad (p = 1, 2, \dots, M) \quad (5)$$

where  $P_{ia}^{ia} \equiv \partial P_{ia} / \partial \tau_p$ .

## 2. Fisher Information Matrix

The Fisher information matrix on the right of (2) now has as a typical element

$$I_{pq} \equiv E \left( \frac{\partial \ell}{\partial \tau_p} \frac{\partial \ell}{\partial \tau_q} \right) = \sum_{i=1}^n \sum_{j=1}^n \sum_{a=1}^N \sum_{b=1}^N \frac{P_{ia}^{ia} P_{jb}^{jb}}{P_{ia} Q_{ia} P_{jb} Q_{jb}} \text{Cov}(u_{ia}, u_{jb})$$

(  $p, q = 1, 2, \dots, M$  )

Because of local independence and random sampling of examinees,

$$\text{Cov}(u_{ia}, u_{jb}) = \delta_{ij} \delta_{ab} P_{ia} Q_{ia}$$

where  $\delta_{st} = 1$  if  $s = t$ ,  $\delta_{st} = 0$  otherwise. Thus the typical element is

$$I_{pq} = \sum_{i=1}^n \sum_{a=1}^N \frac{P_{ia}^{ia} P_{ia}^{ia}}{P_{ia} Q_{ia}} \quad (p, q = 1, 2, \dots, M) \quad (6)$$

Note that  $p_{ia}$  is zero unless either  $p$  and  $a$  refer to the same person, or  $p$  and  $i$  refer to the same item. Thus

$$\|I_{pq}\| = \begin{bmatrix} S_1 & 0 & \dots & 0 & f_{11} & f_{12} & \dots & f_{1N'} \\ 0 & S_2 & \dots & 0 & f_{21} & f_{22} & \dots & f_{2N'} \\ \vdots & \vdots & & \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & S_n & f_{n1} & f_{n2} & \dots & f_{nN'} \\ \hline f'_{11} & f'_{12} & \dots & f'_{1n} & t_1 & 0 & \dots & 0 \\ f'_{21} & f'_{22} & \dots & f'_{2n} & 0 & t_2 & \dots & 0 \\ \vdots & \vdots & & \vdots & \vdots & \vdots & & \vdots \\ f'_{N'1} & f'_{N'2} & \dots & f'_{N'n} & 0 & 0 & \dots & t_{N'} \end{bmatrix} \quad (7)$$

where  $N' \equiv N - 2$ ,  $S_i$  is the 3-by-3 Fisher information matrix for  $a_i$ ,  $b_i$ , and  $c_i$ ,  $t_a$  is the Fisher information for examinee  $a$ , and  $f_{ia}$  is the 3-by-1 joint Fisher information vector for item  $i$  and examinee  $a$ :

$$f_{ia} = \frac{\partial P_{ia} / \partial \theta_a}{P_{ia} Q_{ia}} \begin{bmatrix} \partial P_{ia} / \partial a_i \\ \partial P_{ia} / \partial b_i \\ \partial P_{ia} / \partial c_i \end{bmatrix}$$

### 3. Matrix Inversion

The following general formula for inverting a partitioned matrix may be applied to (7)

$$\begin{bmatrix} S & F \\ F' & T \end{bmatrix}^{-1} \equiv \begin{bmatrix} S^{-1} + S^{-1} F Z^{-1} F' S^{-1} & -S^{-1} F Z^{-1} \\ -Z^{-1} F' S^{-1} & Z^{-1} \end{bmatrix} \quad (8)$$

where

$$Z \equiv T - F' S^{-1} F \quad (9)$$

The matrix  $S$  is easily inverted since it is a diagonal supermatrix:

$$S^{-1} = \begin{bmatrix} S_1^{-1} & 0 \\ 0 & \end{bmatrix}$$

The notation on the right denotes a diagonal matrix with diagonal elements  $S_i^{-1}$ . These last are easily computed since each  $S_i$  is only 3 by 3.

All the matrix operations indicated on the right side of (8) can be carried out on the computer without difficulty, with one exception: the inversion of  $Z$ , which is  $N'$  by  $N'$ . The approximation used here to invert  $Z$  relies on grouping the  $\theta_a$  into 16 class intervals of width 0.5, covering the range  $-5 \leq \theta_a \leq 3$ . Each  $\theta_a$  in a given class interval is replaced by the midpoint of the interval.

Now  $T$  will be a diagonal supermatrix  $T \equiv \begin{bmatrix} & & 0 \\ & T_g & \\ 0 & & \end{bmatrix}$ , where  $T_g \equiv t_g I$  is a scalar matrix with dimensions  $N_g$  by  $N_g$ , and  $N_g$  is the number of people in class interval  $g$ . Also,  $F$  will be a row vector of 16 matrices, the columns of any one matrix being all identical:

$$F \equiv \{f_{11}^{1'}, f_{22}^{1'}, \dots, f_{1616}^{1'}\} \quad (10)$$

where  $f_g \equiv \{f_{ia}\}$  for any examinee  $a$  in class interval  $g$  and  $1_g$  is a unit vector whose length is  $N_g$ .

The product  $F'S^{-1}F$  can now be written as a 16-by-16 supermatrix:

$$F'S^{-1}F = \begin{bmatrix} 1_g' f_g' S^{-1} f_g 1_g \\ \vdots \end{bmatrix}$$

Denote the scalar  $f_g' S^{-1} f_h$  by  $w_{gh}$ . We now have

$$Z \equiv T - \begin{bmatrix} M_{gh} \end{bmatrix} \quad (11)$$

$$M_{gh} \equiv w_{gh} 1_g 1_h' \quad (12)$$

For computation purposes,  $Z$  still has  $N'$  rows and columns, not just 16. For the usual sample size, it is still not feasible to invert  $Z$  with a standard inversion program.

Consider the problem of inverting  $Z_{11}$ , the  $N_1$ -by- $N_1$  upper left corner of  $Z$ . By (11), (12), and a standard formula,

$$Z_{11}^{-1} = [T_1 - w_{11} \begin{smallmatrix} 1 & 1' \\ 1 & 1 \end{smallmatrix}]^{-1} = T_1^{-1} + \frac{w_{11} T_1^{-1} \begin{smallmatrix} 1 & 1' \\ 1 & 1 \end{smallmatrix} T_1^{-1}}{1 - w_{11} \begin{smallmatrix} 1 & 1' \\ 1 & 1 \end{smallmatrix} T_1^{-1} \begin{smallmatrix} 1 \\ 1 \end{smallmatrix}} \quad (13)$$

Since  $T_1 \equiv t_1 I$ , where  $t_1$  is scalar, this becomes

$$Z_{11}^{-1} = \frac{I}{t_1} + \frac{w_{11} \begin{smallmatrix} 1 & 1' \\ 1 & 1 \end{smallmatrix}}{t_1^2 - t_1 w_{11} N_1}$$

Next, the upper left 2-by-2 supermatrix in  $Z$  can be inverted as in (8), using the standard formula for the inversion of a partitioned matrix:

$$\begin{bmatrix} Z_{11} & Z_{12} \\ Z_{21} & Z_{22} \end{bmatrix}^{-1} = \left[ \begin{array}{c|c} Z_{11}^{-1} + Z_{11}^{-1} Z_{12} H^{-1} Z_{21} Z_{11}^{-1} & -Z_{11}^{-1} Z_{12} H^{-1} \\ \hline -H^{-1} Z_{21} Z_{11}^{-1} & H^{-1} \end{array} \right] \quad (14)$$

where  $H = Z_{22} - Z_{21} Z_{11}^{-1} Z_{12}$ . It can be seen that  $H$  has the same general form as  $Z_{11}$  and can thus be inverted as in (13); so (14) can readily be calculated.

Next, substitute (14) for  $Z_{11}^{-1}$  in the foregoing procedure, and repeat this procedure, in such a way as to invert the upper

left 3-by-3 supermatrix in  $Z$ . A total of fifteen repetitions enable us to invert the 16-by-16 supermatrix  $Z$ . Equation (8) is now used for one final inversion, the result being the desired variance-covariance matrix of all  $N + 3n - 2$  parameters.

The 16-by-16 variance-covariance supermatrix for the  $\hat{\theta}_a$  consists of 256 blocks. The elements are all the same within a block except for diagonal blocks, each of which has a variance (instead of a covariance) repeated along its diagonal. Any two examinees in the same class interval will have identical  $\text{Var } \hat{\theta}$  and identical sampling covariances with any other given parameter estimate.

#### 4. Reparameterization

In Section 1, in order to have identifiable parameters, an origin and scale was chosen so that  $\theta_{N-1}$  and  $\theta_N$  had arbitrary preassigned values. Any other choice of origin and scale would result in a linear transformation of parameters. The likelihood function would remain unchanged for every pattern of item responses.

The choice of unit (but not the choice of origin) has one completely obvious effect on the sampling errors of parameter estimates. If the unit is changed, the standard errors for the  $\hat{b}$ 's and  $\hat{\theta}$ 's will be multiplied by the ratio of the new scale unit to the old scale unit. The standard errors for the  $\hat{a}$ 's will be divided by this ratio. A second important effect is easily overlooked: the standard error

of the maximum likelihood estimator depends not only on the choice of scale, but also on how the (origin and) scale is specified.

Suppose that the true numerical values of all  $\theta_a$  ( $a = 1, \dots, N$ ) are specified on some arbitrary scale. Suppose next that our test is too difficult for examinee  $N$ . This means that the likelihood function is rather insensitive to variations in  $\theta_N$ . If we could repeat our testing with several parallel test forms, we would find a wide range of estimates of  $\theta_N$ . In such a situation, the difference between true  $\theta_{N-1}$  and  $\theta_N$  clearly cannot be estimated well from the examinee responses. If we define the scale by treating  $\theta_N$  and  $\theta_{N-1}$  as known, our estimates of every  $\theta_a$  may fluctuate grossly, simply because the scale unit  $\theta_N - \theta_{N-1}$  is not well determined by the data.

Suppose next that we relabel all examinees so that examinees  $N - 1$  and  $N$  are not the same examinees as before. The ability scale has not been changed from the preceding paragraph; it is the procedure for defining the scale that has been changed. The true  $\theta$  for each examinee is still the same as before. Suppose the new examinees  $N - 1$  and  $N$  are both at ability levels where our test measures accurately. If, further, the true  $\theta_{N-1}$  and  $\theta_N$  are substantially different from each other, the difficulty of the previous paragraph disappears: Throughout the ability range where the test is designed to measure accurately, the standard errors of all  $\hat{\theta}_a$  may be reasonably small.



For example, suppose on some scale  $\theta_1 = -3$ ,  $\theta_2 = -2$ ,  $\theta_3 = -1$ ,  $\theta_4 = 0$ ,  $\theta_5 = 1$ ,  $\theta_6 = 2$ ,  $\theta_7 = 3$ . We can specify this same scale in terms of any two of these  $\theta$ 's. The standard errors that we obtain will depend in an overwhelming way not just on the ability scale, but on how we specify it. We cannot rectify the standard errors by some simple procedure, such as multiplying each by a constant.

For this reason, our procedure for specifying the ability scale should depend only on parameters or functions of parameters that are accurately determined by the data. A robust mean of the  $\theta_a$  might seem attractive; however, any function of the  $\theta_a$  is counterindicated by the fact that sometimes  $\hat{\theta}_a = \pm \infty$ .

The procedure used here is to choose a set of  $m$  discriminating, moderately easy items and a set of  $r$  discriminating, moderately hard items. We will hereafter define the origin and unit for our new parameters, to be denoted by capital letters, so that the mean of the (true)  $B$ -parameters for the easy items is zero, and the mean for the hard items is one.

Our new parameters are related to our old parameters (from Section 2 or from Section 5) by linear transformations:

$$A_i \equiv ka_i, \quad B_i \equiv K + b_i/k, \quad C_i \equiv c_i, \quad \theta_a \equiv K + \theta_a/k, \quad (15)$$

$$(a = 1, 2, \dots, N; \quad i = 1, 2, \dots, n)$$

where  $k$  and  $K$  are transformation constants to be determined.

Since

$$\bar{B}_0 \equiv \frac{1}{m} \sum B_i = 0, \quad \bar{B}_1 \equiv \frac{1}{r} \sum B_i = 1, \quad (16)$$

the values of  $k$  and  $K$  are found by substituting (15) into (16) and solving for  $k$  and  $K$  :

$$k = \bar{b}_1 - \bar{b}_0, \quad K = -\frac{\bar{b}_0}{k}, \quad (17)$$

where  $\bar{b}_0$  and  $\bar{b}_1$  are means for  $m$  and  $r$  items, respectively.

To find the variance-covariance matrix for estimates of the upper-case parameters, rewrite (15) as

$$\theta_a = (\theta_a - \bar{b}_0)/k, \quad A_i = ka_i, \quad B_i = (b_i - \bar{b}_0)/k, \quad (18)$$

$$C_i = c_i.$$

Because of the special properties of maximum likelihood estimators, equations (18) still hold when estimators are substituted for parameters.

Thus the sampling variances and covariances for estimates of the new parameters can be computed from the sampling variances and covariances already obtained at the end of Section 3. Formulas for doing this can be written down from (18) by using the 'delta' method (Kendall & Stuart, 1969, Chapter 10). For example,

$$\begin{aligned}
 \text{Cov}(\hat{A}_i, \hat{\theta}_a) &= \text{Cov}(\hat{a}_i, \hat{\theta}_a) - \text{Cov}(\hat{a}_i, \hat{b}_0) \left[ \frac{\theta_a - \bar{b}_0}{k} \text{Cov}(\hat{a}_i, \hat{k}) \right. \\
 &\quad \left. + \frac{a_i}{k} \text{Cov}(\hat{\theta}_a, \hat{k}) - \frac{a_i}{k} \text{Cov}(\hat{b}_0, \hat{k}) - \frac{a_i(\theta_a - \bar{b}_0)}{k^2} \text{Var } \hat{k} \right], \quad (19) \\
 \text{Cov}(\hat{b}_0, \hat{k}) &= \text{Cov}(\hat{b}_1, \hat{b}_0) - \text{Var } \hat{b}_0, \\
 \text{Cov}(\hat{b}_1, \hat{b}_0) &= \frac{1}{mr} \sum \sum^{m,r} \text{Cov}(\hat{b}_i, \hat{b}_j).
 \end{aligned}$$

### 5. Parameter Estimation

The maximum likelihood estimators (MLE) satisfy the likelihood equations (5). In (5), there is one equation for each parameter omitting  $\theta_{N-1}$  and  $\theta_N$ . If all  $N + 3n \equiv M + 2$  MLE are linearly transformed, as for example in (15), the transformed parameters will still satisfy the likelihood equations.

Since the origin and scale for the new parameters is chosen to satisfy (16), then the appropriate  $k$  and  $K$  are obtained from (17) after replacing  $\bar{b}_0$  and  $\bar{b}_1$  by their MLE. The likelihood function (3) is unaffected by these linear transformations.

The computer program LOGIST identifies the parameters by still another choice of origin and scale:

1. a certain truncated mean of the  $\hat{\theta}_a$  ( $a = 1, 2, \dots, N$ ) is set equal to zero,
2. a certain truncated standard deviation of the  $\hat{\theta}_a$  is set equal to one.

We will use the usual lower case symbols for parameters on this LOGIST scale. This should not cause confusion, since the lower-case parameters of Sections 1-3 will not be needed again.

If we start with LOGIST  $\hat{a}_i$ ,  $\hat{b}_i$ ,  $\hat{c}_i$ , and  $\hat{\theta}_a$  and determine  $k$  and  $K$  so that  $\hat{B}_0 = 0$  and  $\hat{B}_1 = 1$ , then the  $\hat{A}_i$ ,  $\hat{B}_i$ ,  $\hat{C}_i$  ( $i = 1, 2, \dots, n$ ), and the  $\hat{\theta}_a$  ( $a = 1, 2, \dots, N$ ), calculated by substituting estimated values into (15), will still satisfy the likelihood equations. The upper-case parameter estimates so obtained should have the sampling variance-covariance matrix found theoretically at the end of Section 4. Our remaining task is to compare an empirically determined variance-covariance matrix of MLE's with the corresponding theoretical matrix.

## 6. Recapitulation

We have used, at different points, three different arbitrary scales for our parameters:

1.  $\theta_N$  and  $\theta_{N-1}$  are assigned arbitrarily.
2. The origin is set at  $\bar{B}_0$ , the unit is  $\bar{B}_1$ .

3. The origin is set at a truncated mean of the  $\theta_a$ ,  
the unit is a truncated standard deviation of the  
 $\theta_a$ .

Scale 1 (denoted by lower-case symbols) is most convenient mathematically for the difficult task of inverting the  $M$ -by- $M$  information matrix. Scale 1 is not useful for practical purposes, however, since its use grossly inflates all the sampling variances.

Scale 2 (denoted by upper-case symbols) seems the simplest choice in an attempt to keep the sampling error in the estimated origin and unit as small as possible. The sampling variances computed for scale 1 are transformed (see eq. 19) to values appropriate for scale 2. Although scale 2 is not the familiar one, the two item sets used to specify the scale can be chosen so that the numerical values of  $\hat{A}_i$ ,  $\hat{B}_i$ ,  $\hat{C}_i$  differ little from the familiar  $\hat{a}_i$ ,  $\hat{b}_i$ , and  $\hat{c}_i$  produced by LOGIST.

Scale 3 (hereafter denoted by lower-case symbols) is the scale used by LOGIST.

## 7. Empirical Estimation Procedures

As already stated, our theoretical results can be trusted only if they are shown to be in reasonable agreement with empirical results. For this purpose, artificial data  $\|u_{ia}\|$  were created representing the administration of a 45-item test to a random sample of 1500

examinees. The 1500  $\theta_a$  were a spaced sample drawn from a distribution of abilities from a regular test administration. Six replicate matrices of  $||u_{ia}||$  were independently generated, using the same item parameters and the same 1500  $\theta_a$ . The variation in responses across these matrices thus represents random fluctuations in  $u_{ia}$  for fixed  $a_i$ ,  $b_i$ ,  $c_i$  and  $\theta_a$ .

Further replication was also built in: items 16-30 and items 31-45 had the same item parameters as items 1-15. The true lower-case and upper-case item parameters are shown in Table 1 for items 1-15.

Six independent runs were made on LOGIST, one for each group of 1500 examinees. For each run separately,  $\hat{b}_0$  was calculated from items 4-9, 19-24, 34-39;  $\hat{b}_1$  was calculated from items 10-15, 25-30, 40-45. It is convenient for our ultimate interpretation of the standard errors to be obtained that the true  $\bar{b}_1 - \bar{b}_0 = .671 - (-.305) = .976$ . Since this is close to 1.0, the scale unit for the capitalized parameters is very close to the scale unit for the lower-case (LOGIST) parameters.

For each run separately, all lower-case parameter estimates were linearly transformed as in (15) to the upper-case scale, using estimated  $k$  and  $K$  values. For the data reported in subsequent sections, the true  $k = .976$  and the true  $K = .312$ . Since the six runs are independent, an unbiased empirical estimate of the sampling variance of any parameter estimate  $T$  is given by

Table 1

True (Upper Case) Item Parameters

Item No.	<u>A</u>	<u>a</u>	<u>B</u>	<u>b</u>	<u>C</u> or <u>c</u>
1	.96	.99	-1.75	-2.01	.17
2	.34	.35	-1.33	-1.61	.17
3	1.34	1.38	-.80	-1.09	.17
4	.76	.78	-.48	-.77	.17
5	.41	.42	-.38	-.67	.17
6	.90	.92	-.04	-.34	.17
7	.90	.92	.16	-.15	.17
8	1.04	1.06	.31	.00	.17
9	1.31	1.34	.42	.11	.13
10	1.46	1.50	.58	.26	.34
11	.85	.87	.79	.46	.17
12	.60	.62	.90	.57	.17
13	1.06	1.09	1.01	.68	.25
14	1.36	1.39	1.23	.90	.29
15	1.46	1.50	1.50	1.16	.18

$$s_T^2 = \frac{6}{5} \left[ \frac{1}{6} \sum \hat{T}^2 - \left( \frac{1}{6} \sum \hat{T} \right)^2 \right] \quad (20)$$

the sum being across the six LOGIST runs. If the  $\hat{T}$  in (20) were normally distributed,  $s_T^2/\sigma_T^2$  would have an F distribution with 5 and  $\infty$  degrees of freedom.

Since three different items have identical item parameters, the  $s_T^2$  for a single item parameter can be averaged across these three items to yield the best available unbiased estimate:

$$\bar{s}_T^2 = \frac{1}{3} \sum s_T^2 \quad (21)$$

Note that it would be incorrect to pool all 18 values of  $\hat{T}$  in an equation like (20), since  $\hat{T}$  from the same LOGIST run are not independent.

If  $T_i$  and  $S_i$  represent two different item parameters in the same item

$$\bar{s}(\hat{T}_i, \hat{S}_i) = \frac{1}{3} \sum s(\hat{T}_i, \hat{S}_i) \quad (22)$$

which is the same as (21) except that covariances are substituted for variances. If  $\hat{T}_i$  and  $\hat{S}_j$  represent item parameters in different items, then there are nine different sample covariances to be summed:



$$\bar{s}(\hat{T}_i, \hat{S}_j) = \frac{1}{9} \sum \sum s(\hat{T}_i, \hat{S}_j) \quad (23)$$

If  $T$  is an ability parameter, (20) still holds. For our purposes, replacing  $T$  by  $\theta$ , we can write

$$\bar{s}_{\theta}^2 = \frac{1}{N_g} \sum s_{\theta}^2 \quad (24)$$

where the sum is over all examinees in group  $g$ . When  $\theta$  is at the midpoint of interval  $g$ , this average should be roughly equal to the  $\sigma_{\theta}$  obtained in Section 4.

If subscripts  $a$  and  $b$  denote different examinees in group  $g$ ,

$$\bar{s}(\hat{\theta}_a, \hat{\theta}_b) = \frac{2}{N_g(N_g - 1)} \sum_{a>b} s(\hat{\theta}_a, \hat{\theta}_b) \quad (25)$$

where the sum is over all pairs of examinees in group  $g$ . If  $a$  and  $b$  denote examinees in groups  $g$  and  $h$  respectively ( $g \neq h$ ), then

$$\bar{s}(\hat{\theta}_a, \hat{\theta}_b) = \frac{1}{N_g N_h} \sum_{a=1}^{N_g} \sum_{b=1}^{N_h} s(\hat{\theta}_a, \hat{\theta}_b) \quad (26)$$

Finally, if  $T_1$  is an item parameter and examinee  $a$  is in group  $g$ , then

$$\bar{s}(\hat{T}_i, \hat{\theta}_a) = \frac{1}{3N_g} \sum_g \sum_s s(\hat{T}_i, \hat{\theta}_a) \quad (27)$$

In computing (24) - (27), examinees are grouped on their true values, not on their estimated values.

A problem arises when an examinee obtains a perfect score or a zero score. In this case his  $\hat{\theta}$  is infinite and cannot be advantageously used. Instead of making some ad hoc adjustment, the 17 examinees for whom this occurred were simply removed from the group of examinees studied, leaving  $N = 1483$ . This has the effect of slightly biasing  $\bar{s}_0$  for the remaining most extreme  $\theta$  values.

#### 8. Numerical Standard Errors

Since the  $c$  parameter of an easy item usually cannot be accurately estimated, LOGIST in ordinary use does not estimate them individually. This would prevent the empirical standard errors of Section 7 from agreeing with the theoretical standard errors of Section 4. Since our main purpose is to show that the method of Section 4 can give useful results, the empirical and theoretical standard errors reported here are all estimated or calculated under the condition that the true values of  $c_i$  are known for  $i = 1, 2, 3, 4, 5, 12$ . Items 1-5 are easy items, item 12 was included because of its low  $a_i$ . For empirical work, the true  $c$  values were supplied to LOGIST, which held them fixed while estimating all other parameters. For theoretical work, the rows and columns of (7) corresponding to  $c_1, c_2, c_3, c_4, c_5$ ,

and  $c_{12}$  were simply deleted from the information matrix (7) before inversion.

Table 2 compares the empirical standard errors of Section 7 for  $\hat{B}$  with the theoretical standard errors of Section 4. The last three columns show the squared ratios for the three replications of each item; each of these ratios will have an  $F$  distribution with 5 and  $\infty$  degrees of freedom provided i)  $\hat{B}$  has a normal sampling distribution, ii)  $\hat{B}$  is unbiased, and iii) the theoretical  $\sigma_{\hat{B}}$  from Section 4 is correct. An  $F$  above 2.21 or below .229 is significant at the (two-tailed) 10 percent level. Eleven of the ratios are significant. The number of ratios less than 1 is approximately the same as the number of ratios greater than 1.

In the past, the only available standard errors for item parameters assumed that the  $\theta$  were known. Such standard errors for  $\hat{B}$ , for known  $\theta$ , are given in the second column of the table. A comparison of second and third columns shows very close agreement except for the three easiest items (1,2,3). For these three items, our new theoretical value is larger and agrees better with the empirical value. This gives support to the new theoretical values. The fact that the empirical values (from Section 7) tend to be larger than the theoretical (from Section 4) could be due to  $n$  and  $N$  not being large enough for asymptotic results. A second likely explanation is that LOGIST was not really run to complete convergence.

Table 3 makes comparisons for  $\hat{A}$ . Again the standard errors of  $\hat{A}$  with  $\theta$  unknown agree closely with the results when  $\theta$  is known. The empirical standard errors, although correlating well with the theoretical, seem to be larger. Eleven of the  $F$  ratios are

Table 2  
Theoretical and Empirical Standard Errors for  $\hat{B}$

Item No.	$\sigma_{\hat{B} \theta}$ ( $\theta$ known)	$\sigma_{\hat{B}}$ (Sect. 4)	$\bar{s}_{\hat{B}}$ (Sect. 7)	$s_{\hat{B}}^2/\sigma_{\hat{B}}^2$		
1*	.110	.156	.183	.23	.56	3.34†
2*	.186	.201	.237	1.76	1.49	.93
3*	.045	.071	.063	1.38	.59	.41
4*	.060	.068	.066	.90	.76	1.17
5*	.100	.099	.103	.37	.40	2.48†
6	.125	.121	.131	.28	.63	2.63†
7	.113	.110	.100	1.24	.65	.58
8	.084	.083	.088	2.31†	.97	.16†
9	.055	.055	.067	.37	2.63†	1.47
10	.069	.069	.106	3.19†	3.62†	.33
11	.100	.097	.122	1.45	2.55†	.70
12*	.094	.091	.087	.85	1.27	.66
13	.086	.083	.094	1.01	1.20	1.57
14	.077	.076	.111	1.19	1.49	3.75†
15	.072	.075	.093	.40	2.62†	1.65

† Significant at 10 percent level.

\*The C parameter for these items is treated as known.

Table 3  
Theoretical and Empirical Standard Errors for  $\hat{A}$

Item No.	$\sigma_{\hat{A} \theta}$	$\sigma_{\hat{A}}$	$\bar{s}_{\hat{A}}$	$\frac{s_{\hat{A}}^2}{\sigma_{\hat{A}}^2}$		
1*	.088	.105	.141	.95	.91	3.60†
2*	.044	.046	.039	.88	.51	.74
3*	.097	.117	.094	1.39	.32	.22†
4*	.060	.065	.080	.89	2.77†	.86
5*	.045	.047	.054	.63	2.44†	.93
6	.103	.102	.123	1.54	.30	2.51†
7	.105	.105	.147	1.30	2.25†	2.35†
8	.113	.115	.159	1.29	3.20†	1.29
9	.123	.128	.182	1.89	3.39†	.80
10	.184	.193	.160	.71	.55	.79
11	.115	.120	.132	1.42	1.85	.34
12*	.060	.060	.076	.95	2.94†	.94
13	.151	.157	.187	2.40†	1.08	.79
14	.209	.218	.240	1.32	.91	1.43
15	.222	.233	.182	.25	.65	.93

†Significant at 10 percent level.

\*The C parameter for these items is treated as known.

significant. Similar statements apply to Table 4, which shows the comparisons for  $\hat{C}$ .

Table 5 compares standard errors for  $\hat{\theta}$ . Let us leave column 3 for later discussion. Columns 4 and 5 show standard errors of  $\hat{\theta}$  corresponding to the  $\theta$  value in the first column; column 6, however, is computed from (2) for the group of  $N_g$  people falling in the class interval with midpoint  $\theta$ . There is good agreement between empirical and theoretical standard errors except for  $\theta < -1.5$ . For low  $\theta$ , asymptotic results do not appear with the usual  $n$  and  $N$ .

Table 5 shows close agreement of our standard error from Sections 2-4 with the standard error of  $\hat{\theta}$  when the item parameters are known. The agreement shown here and in previous tables suggests that (1) is a good approximation to the diagonal of (2) and similarly for item parameters, that (2) agrees well with the empirical standard errors.

A comparison of the third and fifth columns in Table 5 shows what happens to  $\sigma_{\hat{\theta}}$  when all  $C_i$  must be estimated from the data: For  $\theta < -1$ ,  $\sigma_{\hat{\theta}}$  is sharply affected; for  $0 < \theta < 2.5$ , there is very little effect.

Table 6 contains the squared ratios of the empirical standard errors to the theoretical standard errors for the five  $\theta$  closest to the midpoint of the intervals, and within at least .1 of the midpoint. Two of the groups had only two abilities within this restriction. If similar caveats apply as for the item parameters these ratios will have an  $F$  distribution with five and  $\infty$  degrees of freedom. Only eight of the ratios are significant at the two-tailed 10% level, and only 16 are greater than 1.

Table 4

Theoretical and Empirical Standard Errors in  $\hat{C}$

Item No.*	$\sigma_{\hat{C} \theta}$	$\sigma_{\hat{C}}$	$\bar{s}_{\hat{C}}$	$s_{\hat{C}}^2/\sigma_{\hat{C}}^2$		
6	.056	.058	.063	.39	.44	2.79†
7	.049	.050	.038	.40	.35	.95
8	.037	.037	.045	3.08†	.76	.43
9	.024	.025	.039	.80	4.71†	1.83
10	.025	.026	.034	2.24†	2.68†	.27
11	.036	.037	.043	.98	2.67†	.41
13	.026	.027	.037	.89	1.88	2.90†
14	.019	.020	.028	2.98†	2.55†	.43
15	.015	.015	.016	.64	1.23	1.71

†Significant at 10 percent level.

\* $C_1, \dots, C_5$ , and  $C_{12}$  are treated as known.

Table 5  
Theoretical and Empirical Standard Errors for  $\hat{\theta}$

$\theta$	$N_g$	All $C_i$ unknown	$C_1$ to $C_5$ and $C_{12}$ treated as known		
		$\sigma_{\hat{\theta}}$	$\sigma_{\hat{\theta} A,B,C}$	$\sigma_{\hat{\theta}}$	$\bar{s}_{\hat{\theta}}$
-2.75	10	2.090	.951	.966	*
-2.25	35	1.296	.686	.699	1.134
-1.75	93	.861	.516	.525	.797
-1.25	219	.607	.400	.404	.427
-.75	332	.456	.341	.342	.332
-.25	326	.349	.295	.295	.279
.25	227	.278	.262	.263	.274
.75	136	.261	.260	.261	.286
1.25	77	.303	.289	.290	.349
1.75	25	.422	.384	.387	.412
2.25	3	.628	.575	.580	*
2.75	0	.931	.874	.878	*

\*Not computed because of small  $N_g$ .



Table 6  
F Ratios for  $\hat{\theta}$

$\theta$	$s_{\hat{\theta}}^2 / \sigma_{\hat{\theta}}^2$				
-2.75	3.73†	4.41†			
-2.25	.85	.78	.43	11.34†	1.16
-1.75	.57	1.90	1.62	.32	18.95†
-1.25	.98	.63	.96	.95	.77
-.75	.26	.94	.63	.81	.63
-.25	.71	1.81	.73	.04†	.48
.25	.18†	.98	.74	.80	.77
.75	.61	.35	1.41	1.21	.64
1.25	2.76†	1.82	.98	1.08	1.84
1.75	.67	.41	1.08	1.45	1.78
2.25	.11†	.36			
2.75*					

†Significant at 10 percent level.

\*There were no  $\theta$  between 2.65 and 2.85.

Table 7 presents the theoretical standard errors of  $\hat{A}$ ,  $\hat{B}$ , and  $\hat{C}$ , obtained by the method of Sections 2-4, when all  $C_i$  must be estimated from the data. It is interesting to compare these values with those in Tables 2-4 where  $C_1, \dots, C_5$ , and  $C_{12}$  were treated as known. We find that the standard errors of  $\hat{B}_1$  to  $\hat{B}_5$  are increased drastically by ignorance of  $C_1$  to  $C_5$ ; all other  $\sigma(\hat{B}_i)$  are much increased, except for  $i = 11, 13$ , and  $14$ . All  $\hat{A}_i$  show sharply increased standard errors. For items for which  $C_i$  must be estimated, on the other hand, the standard errors of  $\hat{C}_i$  are little affected by knowledge or ignorance of  $C_1, \dots, C_5, C_{12}$ . A likely explanation for this is that errors in estimating the scale unit  $B_1$  affect the standard errors of the  $\hat{A}_i$  and the  $\hat{B}_i$ , but not of the  $\hat{C}_i$ .

We have found in Tables 2-7 some illustrative answers to the question: How do estimation errors on one set of items affect the accuracy of estimated parameters for a different set of items? Such effects could not be quantified until now since the standard error of an item parameter estimate was previously known only for fixed  $\theta$ . It is only through the sampling fluctuations of  $\hat{\theta}$  that estimation errors for one item can affect parameter estimates for another item.

With 18  $C_i$  treated as known, the Fisher information matrix inverted for this study has  $3 \times 45 - 18 = 1497$  rows and columns. The matrix inversion by the method of Section 4 used 1232K bytes of memory on an IBM 3031 and took 32 seconds. The computer program dealt with a 45-item test; it did not take advantage of the fact that the 45 items consisted of 3 replicate sets of 15 items each.

Table 7

Standard Errors (2) of Item Parameters when  
All  $C_i$  Must be Estimated

Item No.	$\sigma_B$	$\sigma_A$	$\sigma_C$
1	.52	.23	.60
2	.54	.13	.72
3	.35	.32	.10
4	.26	.15	.14
5	.97	.10	.32
6	.19	.18	.07
7	.16	.18	.06
8	.14	.21	.041
9	.12	.26	.026
10	.11	.32	.026
11	.10	.18	.039
12	.18	.14	.07
13	.09	.23	.027
14	.08	.31	.020
15	.10	.33	.015

In order to verify the numerical accuracy of the inversion, the information matrix and the variance-covariance matrix were multiplied. The result was an identity matrix accurate to 10 decimal places. The variance-covariance matrix obtained in double precision agreed with the matrix obtained in quadruple precision to all six decimal places printed.

### 9. Sampling/Covariances and Correlations

When item parameters are known,  $\hat{\theta}_a$  and  $\hat{\theta}_b$  ( $a \neq b$ ) are uncorrelated. When ability parameters are known, estimated item parameters for different items are uncorrelated. When both item and ability parameters are estimated, in general all estimates are correlated. The computer printout of the sampling correlations for the present study consists of 10 correlation matrices. These need only be summarized here.

Table 8 shows the theoretical ( T ) and empirical ( E ) correlations between estimates of two different parameters for the same item. The correlations are generally substantial. For comparison, the theoretical correlations when the abilities are known are included. The empirical correlations are obtained by dividing the estimated sampling covariance by the square roots of the estimated sampling variances. If the empirical correlations here have roughly 15 degrees of freedom, their standard error is roughly  $(1 - \rho^2)/\sqrt{15} = .26(1 - \rho^2)$ . In view of their standard errors, there is very satisfactory agreement of empirical with theoretical correlations.

Table 9 shows both theoretical and empirical correlations for the  $\hat{B}_i$  ( $i = 1, 2, \dots, 15$ ). The corresponding standard errors are

Table 8

Theoretical ( T ) and Empirical ( E ) Sampling Correlations Between  
Two Parameter Estimates for the Same Item

Item	$\rho_{AB}^{\wedge\wedge} \theta$	$\rho_{AB}^{\wedge\wedge}$		$\rho_{BC}^{\wedge\wedge} \theta$	$\rho_{BC}^{\wedge\wedge}$		$\rho_{AC}^{\wedge\wedge} \theta$	$\rho_{AC}^{\wedge\wedge}$	
		T	E		T	E		T	E
1	.82	.86	.87						
2	.80	.82	.88						
3	.55	.70	.65						
4	.42	.52	.76						
5	.35	.38	.44						
6	.73	.70	.53	.92	.90	.92	.76	.70	.53
7	.67	.64	.66	.90	.88	.77	.76	.71	.79
8	.56	.52	.26	.83	.81	.85	.72	.67	.58
9	.37	.33	.50	.69	.67	.87	.65	.60	.81
10	.41	.42	.68	.69	.68	.93	.61	.61	.74
11	.40	.42	.70	.75	.74	.89	.77	.77	.83
12	-.55	-.51	-.79						
13	.22	.21	.06	.60	.59	.66	.69	.70	.67
14	.06	.03	.35	.45	.42	.61	.58	.59	.68
15	-.19	-.25	-.81	.25	.21	-.18	.53	.54	.56

TABLE 9  
EXPERIMENTAL (E) AND THEORETICAL (T) STANDARD ERRORS (DIAGONALS) AND  
CORRELATIONS FOR TRANSFORMED B (DECIMAL POINTS OMITTED)

		B 1	B 2	B 3	B 4	B 5	B 6	B 7	B 8	B 9	B 10	B 11	B 12	B 13	B 14	B 15
B 1	E	(183)	141	284	045	193	-158	-028	-004	014	-122	289	016	-031	-064	-147
	T	(156)	264	509	334	158	-124	-128	-088	-040	034	-001	039	-005	-022	-050
B 2	E	141	(237)	541	286	-092	-036	126	-005	-252	-105	029	-064	-247	360	-040
	T	264	(201)	284	184	078	-066	-069	-046	-017	026	007	022	-002	-018	-039
B 3	E	284	541	(063)	308	040	-091	056	004	-274	-279	268	007	-298	348	-155
	T	509	284	(071)	377	151	-131	-139	-093	-032	048	008	044	-008	-032	-068
B 4	E	045	286	308	(066)	120	-072	-179	038	-362	-308	-007	192	-443	343	218
	T	334	184	377	(068)	066	-130	-130	-089	-040	029	003	028	-005	-018	-039
B 5	E	193	-092	040	120	(103)	-228	-126	-072	046	-205	236	086	-051	-193	126
	T	158	078	151	066	(099)	-117	-113	-088	-062	-004	-009	013	-003	001	002
B 6	E	-158	-036	-091	-072	-228	(131)	014	076	-041	107	-153	002	016	122	-085
	T	-124	-066	-131	-130	-117	(121)	-062	-053	-051	-016	004	-005	001	004	011
B 7	E	-028	126	056	-179	-126	014	(100)	-120	098	121	-050	-221	156	-018	000
	T	-128	-069	-139	-130	-113	-062	(110)	-042	-036	-011	002	-009	002	005	011
B 8	E	-004	-005	004	038	-072	076	-120	(088)	-068	025	-015	101	-062	081	-137
	T	-088	-046	-093	-089	-088	-053	-042	(083)	-007	004	001	-010	002	002	003
B 9	E	014	-252	-274	-362	046	-041	098	-068	(067)	198	037	-129	332	-357	-063
	T	-040	-017	-032	-040	-062	-051	-036	-007	(055)	023	-003	-013	002	000	-005
B 10	E	-122	-105	-279	-308	-205	107	121	025	198	(106)	-193	-137	270	-151	-098
	T	034	026	048	029	-004	-016	-011	004	023	(069)	-035	-052	-043	-062	-087
B 11	E	289	029	268	-007	236	-153	-050	-015	037	-193	(122)	041	-011	-103	-182
	T	-001	007	008	003	-009	004	002	001	-003	-035	(097)	-071	-067	-086	-107
B 12	E	016	-064	007	192	086	002	-221	101	-129	-137	041	(087)	-176	078	005
	T	039	022	044	028	013	-005	-009	-010	-013	-052	-071	(091)	-069	-068	-065
B 13	E	-031	-247	-298	-443	-051	016	156	-062	332	270	-011	-176	(094)	-341	-112
	T	-005	-002	-008	-005	-003	001	002	002	002	-043	-067	-069	(083)	-057	-060
B 14	E	-064	360	348	343	-193	122	-018	081	-357	-151	-103	078	-341	(111)	006
	T	-022	-018	-032	-018	001	004	005	-002	000	-062	-086	-068	-057	(076)	-004
B 15	E	-147	-040	-155	218	126	-085	000	-137	-063	-098	-182	005	-112	006	(093)
	T	-050	-039	-068	-039	002	011	011	003	-005	-087	-107	-065	-060	-004	(075)

given in parentheses in the diagonal. The only theoretical correlations above .20 are among  $\hat{B}_1$ ,  $\hat{B}_2$ ,  $\hat{B}_3$ , and  $\hat{B}_4$ . These are the four easiest items. Any error in estimating the scale unit  $\bar{B}_1 - \bar{B}_0$  would seriously affect all these items in the same way. It is hard to draw other useful generalizations from this table.

The corresponding table for the  $\hat{A}_i$  ( $i = 1, 2, \dots, 15$ ) shows only 3 theoretical correlations above .20:  $\rho_{13} = .27$ ,  $\rho_{14} = .20$ ,  $\rho_{34} = .23$ . With two exceptions ( $\rho_{67} = -.013$ ,  $\rho_{6,12} = -.002$ ), all theoretical correlations are positive.

The highest theoretical correlation among the  $\hat{C}_i$  ( $i = 6, 7, \dots, 11$  and  $13, 14, 15$ ) is  $\rho_{67} = .04$ . All correlations are positive.

The theoretical correlations between  $\hat{A}_i$  and  $\hat{B}_j$  ( $i \neq j$ ) are all below .20 in absolute value, except for items 1-4, which vary from .14 to .38. For  $\hat{B}_i$  and  $\hat{C}_j$  ( $i \neq j$ ;  $j \neq 1, 2, \dots, 5, 12$ ) there are no correlations above .25 in absolute value. For  $\hat{A}_i$  and  $\hat{C}_j$ , there are no correlations above .20 in absolute value.

The theoretical correlations between  $\hat{\theta}_a$  and  $\hat{\theta}_b$  ( $a \neq b$ ) are all less than .04 in absolute value. Between  $\hat{\theta}_a$  and  $\hat{B}_i$ , the largest correlation in absolute value is .15 (when  $i = 1$  and  $\theta = -2.25$ ). Between  $\hat{\theta}_a$  and  $\hat{A}_i$ , the largest is .12 (when  $i = 1$  and  $\theta = -2.25$ ). Between  $\hat{\theta}_a$  and  $\hat{C}_i$ , the largest is .06.

### Summary

When both abilities and item parameters are unknown, the asymptotic sampling variance-covariance matrix developed in this paper appears to provide useful values for the standard errors needed for further research in item response theory. The magnitude of the numerical values in the matrix were very much affected by the method used to define the scale. For a set of artificial data, this variance-covariance matrix compared satisfactorially with empirical results; also with the variance-covariance matrices found by the usual formulas for the case where the abilities are known or where the item parameters are known.

With this matrix, the effect on other items of including items with poorly determined parameters can be studied. Including items with poorly determined  $c$ 's increases the standard errors of all of the  $a$ 's and  $b$ 's but not of the other  $c$ 's. The effect of different distributions of abilities on the accuracy of item parameters can also be studied. Hopefully a goodness-of-fit test can now be developed for the three-parameter model.

The standard errors of item parameters can now be studied for a situation of common occurrence in equating and item banking: Each of two tests containing common items is administered to a different group of examinees; all parameters are estimated in the same LOGIST run. It is of particular interest to determine how the number of common items affects the standard error of the parameter estimates.



References

- Bradley, R. A. & Gart, J. J. The asymptotic properties of ML estimators when sampling from associated populations. Biometrika, 1962, 49, 205-214.
- Haberman, S. J. Maximum likelihood estimates in exponential response models. The Annals of Statistics, 1977, 5, 815-841.
- Kendall, M. G. & Stuart, A. The advanced theory of statistics (Vol. 1, 3rd ed.). New York: Hafner, 1969.
- Lord, F. M. Applications of item response theory to practical testing problems. Hillsdale, N.J.: Lawrence Erlbaum Associates, 1980.

# DISTRIBUTION LIST

## Navy

- |  |  |
|--|--|
| <p>1 Dr. Jack R. Borsting<br/>Provost and Academic Dean<br/>U.S. Naval Postgraduate School<br/>Monterey, CA 93940</p> <p>1 Chief of Naval Education and<br/>Training Liason Office<br/>Air Force Human Resource Laboratory<br/>Flying Training Division<br/>Williams Air Force Base, AZ 85224</p> <p>1 CDR Mike Curran<br/>Office of Naval Research<br/>800 North Quincy Street<br/>Code 270<br/>Arlington, VA 22217</p> <p>1 Dr. Pat Federico<br/>Navy Personnel R &amp; D Center<br/>San Diego, CA 92152</p> <p>1 Mr. Paul Foley<br/>Navy Personnel R &amp; D Center<br/>San Diego, CA 92152</p> <p>1 Dr. John Ford<br/>Navy Personnel R &amp; D Center<br/>San Diego, CA 92152</p> <p>1 Patrick R. Harrison<br/>Psychology Course Director<br/>Leadership and Law Department (7b)<br/>Division of Professional Development<br/>U.S. Naval Academy<br/>Annapolis, MD 21402</p> | <p>1 Dr. Norman J. Kerr<br/>Chief of Naval Technical Training<br/>Naval Air Station Memphis (75)<br/>Millington, TN 38054</p> <p>1 Dr. William L. Maloy<br/>Principal Civilian Advisor for<br/>Education and Training<br/>Naval Training Command, Code 00A<br/>Pensacola, FL 32508</p> <p>1 CAPT Richard L. Martin, USN<br/>Prospective Commanding Officer<br/>USS Carl Vinson (CNV-70)<br/>Newport News Shipbuilding and<br/>Drydock Co.<br/>Newport News, VA 23607</p> <p>1 Dr. James McBride<br/>Navy Personnel R &amp; D Center<br/>San Diego, CA 92152</p> <p>1 Mr. William Nordbrock<br/>Instructional Program Development<br/>Building 90<br/>NET-PDCD<br/>Great Lakes NTC, IL 60088</p> <p>1 Library, Code P201L<br/>Navy Personnel R &amp; D Center<br/>San Diego, CA 92152</p> |
|--|--|

6 Commanding Officer,  
Naval Research Laboratory  
Code 2627  
Washington, DC 20390

1 Psychologist  
ONR Branch Office  
Building 114, Section D  
606 Summer Street  
Boston, MA 02210

1 Office of Naval Research  
Code 437  
800 North Quincy Street  
Arlington, VA 22217

5 Personnel and Training Research  
Programs  
Code 458  
Office of Naval Research  
Arlington, VA 22217

1 Psychologist  
ONR Branch Office  
1030 East Green Street  
Pasadena, CA 91101

1 Office of the Chief of Naval Operations  
Research Development and Studies Branch  
OP-115  
Washington, DC 20350

1 The Principal Deputy Assistant  
Secretary of the Navy (MRA&L)  
4E780, The Pentagon  
Washington, DC 22203

1 Director, Research and  
Analysis Division  
Plans and Policy Department  
Navy Recruiting Command  
4015 Wilson Boulevard  
Arlington, VA 22203

1 Mr. Arnold Rubenstein  
Office of Naval Technology  
800 N. Quincy Street  
Arlington, VA 22217

1 Dr. Worth Scanland, Director  
Research, Development, Test  
and Evaluation  
N-5  
Naval Education and Training Command  
NAS  
Pensacola, FL 32508

1 Dr. Robert G. Smith  
Office of Chief of Naval Operations  
OP-987H  
Washington, DC 20350

1 Dr. Alfred F. Smode  
Training Analysis and Evaluation Group  
Department of the Navy  
Orlando, FL 32813

1 Dr. Richard Sorensen  
Navy Personnel R & D Center  
San Diego, CA 92152

1 Mr. J. B. Sympson  
Naval Personnel R & D Center  
San Diego, CA 92152

1 Dr. Ronald Weitzman  
Code 54 WZ  
Department of Administrative Services  
U.S. Naval Postgraduate School  
Monterey, CA 93940

1 Dr. Robert Wisher  
Code 309  
Navy Personnel R & D Center  
San Diego, CA 92152

1 Dr. Martin F. Wiskoff  
Navy Personnel R & D Center  
San Diego, CA 92152

1 Mr. John H. Wolfe  
Code P310  
U.S. Navy Personnel Research  
and Development Center  
San Diego, CA 92152

1 Mr. Ted M. I. Yellen  
Technical Information Office  
Code 201  
Navy Personnel R & D Center  
San Diego, CA 92152

Army

1 Technical Director  
U.S. Army Research Institute for the  
Behavioral and Social Sciences  
5001 Eisenhower Avenue  
Alexandria, VA 22333

1 Dr. Myron Fischl  
U.S. Army Research Institute for the  
Social and Behavioral Sciences  
5001 Eisenhower Avenue  
Alexandria, VA 22333

1 Dr. Michael Kaplan  
U.S. Army Research Institute  
5001 Eisenhower Avenue  
Alexandria, VA 22333

1 Dr. Milton S. Katz  
Training Technical Area  
U.S. Army Research Institute  
5001 Eisenhower Avenue  
Alexandria, VA 22333

1 Dr. Harold F. O'Neil, Jr.  
Attn: PERI-OK  
Army Research Institute  
5001 Eisenhower Avenue  
Alexandria, VA 22333

1 LTC Michael Plummer  
Chief, Leadership and Organizational  
Effectiveness Division  
Office of the Deputy Chief of Staff  
for Personnel  
Department of the Army  
The Pentagon  
Washington, DC 20301

1 Dr. James L. Raney  
U.S. Army Research Institute  
5001 Eisenhower Avenue  
Alexandria, VA 22333

1 Mr. Robert Ross  
U.S. Army Research Institute for  
the Social and Behavioral Sciences  
5001 Eisenhower Avenue  
Alexandria, VA 22333

1 Dr. Robert Sasmor  
U.S. Army Research Institute for  
the Social and Behavioral Sciences  
5001 Eisenhower Avenue  
Alexandria, VA 22333

1 Commandant  
U.S. Army Institute of Administration  
Attn: Dr. Sherrill  
Ft. Benjamin Harrison, IN 46256

1 Dr. Joseph Ward  
U.S. Army Research Institute  
5001 Eisenhower Avenue  
Alexandria, VA 22333

## Air Force

- 1 Air Force Human Resources Laboratory  
AFHRL/MPD  
Brooks Air Force Base, TX 78235
- 1 U.S. Air Force Office of  
Scientific Research  
Life Sciences Directorate  
Bolling Air Force Base  
Washington, DC 20332
- 1 Dr. Earl A. Alluisi,  
HQ, AFHRL (AFSC)  
Brooks Air Force Base, TX 78235
- 1 Dr. Genevieve Haddad  
Program Manager  
Life Sciences Directorate  
AFOSR  
Bolling Air Force Base  
Washington, DC 20332
- 1 Dr. David R. Hunter  
AFHRL/MOAM  
Brooks Air Force Base, TX 78235
- 1 Research and Measurement Division  
Research Branch, AFMPC/MPCYPR  
Randolph Air Force Base, TX 78148
- 1 Dr. Malcolm Ree  
AFHRL/MP  
Brooks Air Force Base, TX 78235

## Marines

- 1 Dr. H. William Greenup  
Education Advisor (E031)  
Education Center, MCDEC  
Quantico, VA 22134

- 1 Director, Office of Manpower  
Utilization  
HQ, Marine Corps (MPU)  
BCB, Building 2009  
Quantico, VA 22134
- 1 Special Assistant for Marine  
Corps Matters  
Code 100M  
Office of Naval Research  
800 N. Quincy Street  
Arlington, VA 22217
- 1 MAJ Michael L. Patrow, USMC  
Headquarters, Marine Corps  
Code MPI-20  
Washington, DC 20380
- 1 Dr. A. L. Slafkosky  
Scientific Advisor  
Code RD-1  
HQ, U.S. Marine Corps  
Washington, DC 20380

## Coast Guard

- 1 Chief, Psychological Research Branch  
U.S. Coast Guard (G-P-1/2/TP42)  
Washington, DC 20593
- 1 Mr. Thomas A. Warm  
U.S. Coast Guard Institute  
P.O. Substation 18  
Oklahoma City, OK 73169

## Other DoD

- 1 DARPA  
1400 Wilson Boulevard  
Arlington, VA 22209

12 Defense Technical Information Center  
Cameron Station, Building 5  
Attn: TC  
Alexandria, VA 22314

1 Dr. William Graham  
Testing Directorate  
MEPCOM/MEPCT-P  
Ft. Sheridan, IL 60037

1 Director, Research and Data  
OASD (MRA&L)  
3B919, The Pentagon  
Washington, DC 20301

1 Military Assistant for Training  
and Personnel Technology  
Office of the Under Secretary of  
Defense for Research and Engineering  
Room 3D129, The Pentagon  
Washington, DC 20301

1 Dr. Wayne Sellman  
Office of the Assistant Secretary  
of Defense (MRA&L)  
2B269 The Pentagon  
Washington, DC 20301

Civil Government

1 Mr. Richard McKillip  
Personnel R & D Center  
Office of Personnel Management  
1900 E Street, NW  
Washington, DC 20415

1 Dr. Andrew R. Molnar  
Science Education Development  
and Research  
National Science Foundation  
Washington, DC 20550

1 Dr. H. Wallace Sinaiko  
Program Director  
Manpower Research and  
Advisory Services  
Smithsonian Institution  
801 North Pitt Street  
Alexandria, VA 22314

1 Dr. Vern W. Urry  
Personnel R & D Center  
Office of Personnel Management  
1900 E Street, NW  
Washington, DC 20415

1 Dr. Joseph L. Young, Director  
Memory and Cognitive Processes  
National Science Foundation  
Washington, DC 20550

Non-Government

1 Dr. James Algina  
University of Florida  
Gainesville, FL 32611

1 Dr. Erling B. Andersen  
Department of Statistics  
Studiestraede 6  
1455 Copenhagen  
DENMARK

1 Psychological Research Unit  
Department of Defense (Army Office)  
Campbell Park Offices  
Canberra, ACT 2600  
AUSTRALIA

1 Dr. Isaac Bejar  
Educational Testing Service  
Princeton, NJ 08541

1 CAPT J. Jean Belanger  
Training Development Division  
Canadian Forces Training System  
CFTSHQ, CFB Trenton  
Astra, Ontario K0K 1B0  
CANADA

1 Dr. Menucha Birenbaum  
School of Education  
Tel Aviv University  
Tel Aviv, Ramat Aviv 69978  
ISRAEL

1 Dr. Werner Birke  
DezWPs im Streitkrafteamt  
Postfach 20 50 3  
D-5300 Bonn 2  
WEST GERMANY

1 Dr. R. Darrell Bock  
Department of Education  
University of Chicago  
Chicago, IL 60637

1 Liaison Scientists  
Office of Naval Research  
Branch Office, London  
Box 39  
FPO, NY 09510

1 Dr. Robert Brennan  
American College Testing Programs  
P.O. Box 168  
Iowa City, IA 52240

1 Dr. C. Victor Bunderson  
WICAT Inc.  
University Plaza, Suite 10  
1160 S. State Street  
Orem, UT 84057

1 Dr. John B. Carroll  
Psychometric Laboratory  
University of North Carolina  
Davie Hall 013A  
Chapel Hill, NC 27514

1 Charles Myers Library  
Livingstone House  
Livingstone Road  
Stratford  
London E15 2LJ  
ENGLAND

1 Dr. Kenneth E. Clark  
College of Arts and Sciences  
University of Rochester  
River Campus Station  
Rochester, NY 14627

1 Dr. Norman Cliff  
Department of Psychology  
University of Southern California  
University Park  
Los Angeles, CA 90007

1 Dr. William E. Goffman  
Director, Iowa Testing Programs  
334 Lindquist Center  
University of Iowa  
Iowa City, IA 52242

1 Dr. Meredith P. Crawford  
American Psychological Association  
1200 17th Street, N  
Washington, DC 20036

1 Dr. Fritz Drasgow  
Yale School of Organization and  
Management  
Yale University  
Box 1A  
New Haven, CT 06520

1 Dr. Mike Durmeyer  
Instructional Program Development  
Building 90  
NET-PDCD  
Great Lakes NTC, IL 60088

1 ERIC Facility-Acquisitions  
4833 Rugby Avenue  
Bethesda, MD 20014

1 Dr. A. J. Eschenbrenner  
Dept. E422, Bldg. 81  
McDonald Douglas Astronautics Co.  
P.O. Box 516  
St. Louis, MO 63166

1 Dr. Benjamin A. Fairbank, Jr.  
McFann-Gray and Associates, Inc.  
5825 Callaghan  
Suite 225  
San Antonio, TX 78228

1 Dr. Leonard Feldt  
Lindquist Center for Measurement  
University of Iowa  
Iowa City, IA 52242

1 Dr. Richard L. Ferguson  
The American College Testing Program  
P.O. Box 168  
Iowa City, IA 52240

1 Dr. Victor Fields  
Department of Psychology  
Montgomery College  
Rockville, MD 20850

1 Univ. Prof. Dr. Gerhard Fischer  
Psychologisches Institut der  
Universitat Wien  
Liebiggasse 5/3  
A 1010 Wien  
AUSTRIA

1 Prof. Donald Fitzgerald  
University of New England  
Armidale, New South Wales 2351  
AUSTRALIA

1 Dr. Edwin A. Fleishman  
Advanced Research Resources Organization  
Suite 900  
4330 East West Highway  
Washington, DC 20014

1 Dr. John R. Frederiksen  
Bolt, Beranek, and Newman  
50 Moulton Street  
Cambridge, MA 02138

1 Dr. Robert Glaser  
LRDC  
University of Pittsburgh  
3939 O'Hara Street  
Pittsburgh, PA 15213

1 Dr. Daniel Gopher  
Industrial and Management Engineering  
Technion-Israel Institute of  
Technology  
Haifa  
ISRAEL

1 Dr. Bert Green  
Department of Psychology  
Johns Hopkins University  
Charles and 34th Streets  
Baltimore, MD 21218

1 Dr. Ron Hambleton  
School of Education  
University of Massachusetts  
Amherst, MA 01002

1 Dr. Delwyn Harnisch  
University of Illinois  
242b Education  
Urbana, IL 61801

1 Dr. Chester Harris  
School of Education  
University of California  
Santa Barbara, CA 93106

1 Dr. Lloyd Humphreys  
Department of Psychology  
University of Illinois  
Champaign, IL 61820



- 1 Library  
HumRRO/Western Division  
27857 Berwick Drive  
Carmel, CA 93921
- 1 Dr. Steven Hunka  
Department of Education  
University of Alberta  
Edmonton, Alberta  
CANADA
- 1 Dr. Jack Hunter  
2122 Coolidge Street  
Lansing, MI 48906
- 1 Dr. Huynh Huynh  
College of Education  
University of South Carolina  
Columbia, SC 29208
- 1 Prof. John A. Keats  
Department of Psychology  
University of Newcastle  
Newcastle, New South Wales 2308  
AUSTRALIA
- 1 Mr. Jeff Kelety  
Department of Instructional Technology  
University of Southern California  
Los Angeles, CA 90007
- 1 Dr. Michael Levine  
Department of Educational Psychology  
210 Education Building  
University of Illinois  
Champaign, IL 61801
- 1 Dr. Charles Lewis  
Faculteit Sociale Wetenschappen  
Rijksuniversiteit Groningen  
Oude Boteringestraat 23  
9712GC Groningen  
NETHERLANDS
- 1 Dr. Robert Linn  
College of Education  
University of Illinois  
Urbana, IL 61801
- 1 Dr. James Lumsden  
Department of Psychology  
University of Western Australia  
Nedlands, Western Australia 6009  
AUSTRALIA
- 1 Dr. Gary Marco  
Educational Testing Service  
Princeton, NJ 08541
- 1 Dr. Scott Maxwell  
Department of Psychology  
University of Houston  
Houston, TX 77004
- 1 Dr. Samuel T. Mayo  
Loyola University of Chicago  
820 North Michigan Avenue  
Chicago, IL 60611
- 1 Prof. Jason Millman  
Department of Education  
Stone Hall  
Cornell University  
Ithaca, NY 14853
- 1 Dr. Melvin R. Novick  
356 Lindquist Center for Measurement  
University of Iowa  
Iowa City, IA 52242
- 1 Dr. Jesse Orlansky  
Institute for Defense Analyses  
400 Army Navy Drive  
Arlington, VA 22202
- 1 Dr. Wayne M. Patience  
American Council on Education  
GED Testing Service, Suite 20  
One Dupont Circle, NW  
Washington, DC 20036

1 Dr. James A. Paulson  
Portland State University  
P.O. Box 751  
Portland, OR 97207

1 Mr. Luigi Petruccio  
2431 North Edgewood Street  
Arlington, VA 22207

1 Dr. Diane M. Ramsey-Klee  
R-K Research and System Design  
3947 Ridgmont Drive  
Malibu, CA 90265

1 Mr. Minrat M. L. Rauch  
P II 4  
Bundesministerium der Verteidigung  
Postfach 1328  
D-53 Bonn 1  
GERMANY

1 Dr. Mark D. Reckase  
Educational Psychology Department  
University of Missouri-Columbia  
4 Hill Hall  
Columbia, MO 65211

1 Dr. Andrew Rose  
American Institutes for Research  
1055 Thomas Jefferson St., NW  
Washington, DC 20007

1 Dr. Leonard L. Rosenbaum, Chairman  
Department of Psychology  
Montgomery College  
Rockville, MD 20850

1 Dr. Ernst Z. Rothkopf  
Bell Laboratories  
600 Mountain Avenue  
Murray Hill, NJ 07974

1 Dr. Lawrence Rudner  
403 Elm Avenue  
Takoma Park, MD 20012

1 Dr. J. Ryan  
Department of Education  
University of South Carolina  
Columbia, SC 29208

1 Prof. Fumiko Samejima  
Department of Psychology  
University of Tennessee  
Knoxville, TN 37916

1 Dr. Kazuo Shigemasu  
University of Tohoku  
Department of Educational Psychology  
Kawauchi, Sendai 980  
JAPAN

1 Dr. Edwin Shirkey  
Department of Psychology  
University of Central Florida  
Orlando, FL 32816

1 Dr. Robert Smith  
Department of Computer Science  
Rutgers University  
New Brunswick, NJ 08903

1 Dr. Richard Snow  
School of Education  
Stanford University  
Stanford, CA 94305

1 Dr. Robert Sternberg  
Department of Psychology  
Yale University  
Box 11A, Yale Station  
New Haven, CT 06520

1 Dr. Patrick Suppes  
Institute for Mathematical Studies in  
the Social Sciences  
Stanford University  
Stanford, CA 94305

- 1 Dr. Hariharan Swaminathan  
Laboratory of Psychometric and  
Evaluation Research  
School of Education  
University of Massachusetts  
Amherst, MA 01003
- 1 Dr. Kikumi Tatsuoka  
Computer Based Education Research  
Laboratory  
252 Engineering Research Laboratory  
University of Illinois  
Urbana, IL 61801
- 1 Dr. David Thissen  
Department of Psychology  
University of Kansas  
Lawrence, KS 66044
- 1 Dr. Robert Tsutakawa  
Department of Statistics  
University of Missouri  
Columbia, MO 65201
- 1 Dr. Howard Wainer  
Educational Testing Service  
Princeton, NJ 08541
- 1 Dr. David J. Weiss  
N660 Elliott Hall  
University of Minnesota  
75 East River Road  
Minneapolis, MN 55455
- 1 Dr. Susan E. Whitely  
Psychology Department  
University of Kansas  
Lawrence, KS 66044
- 1 Dr. Wolfgang Wildgrube  
Streitkraefteamt  
Box 20 50 03  
D-5300 Bonn 2  
WEST GERMANY